## **Vascular Plants of Williamson County**

*Rhynchosia minima* – LEAST STOUT BEAN [Fabaceae]

Rhynchosia minima (L.) DC., LEAST STOUT BEAN. Perennial vine, twining, not rosetted, several-stemmed at base, growing on support or trailing; shoots with only cauline leaves on flexible stems, puberulent and with scattered, subsessile glandular hairs having yellowish heads. **Stems:** 6-ridged, < 2 mm diameter, with 3 conspicuous ridges descending from each leaf, tough, internodes usually twisted and to 85 mm long, ≥ leaf, glandular hairs absent from ridges, nonglandular hairs downward-pointing with some longer hairs widely spaced along ridges. Leaves: helically alternate, pinnately-3-foliolate, petiolate with pulvinus, with stipules; stipules 2, attached to stem at node, acuminate, 2–3.2 mm long, green to reddish, sometimes with lateral tooth near base, puberulent, deciduous to latedeciduous; petiole pulvinus not noticeably swollen. 1–2.5 mm long, puberulent, above pulvinus strongly 5-ridged, glandular hairs scattered in furrows; rachis with a pair of conspicuous ridges on upper side, like petiole; stipel subtending lateral leaflet 1, subtending terminal leaflet 2, linear extentions of ridges < 0.35 mm long; petiolule = pulvinus ca. 1 mm long, puberulent with glandular hairs; bladelets of leaflets obovate to  $\pm$ diamond-shaped, in range to 30 × 30 mm, broadly tapered to subtruncate or roundish, entire or with 2 broad, rounded lobes above midblade, acute to obtuse at tip, conspicuously 3-veined at base each pinnately veined with principal veins raised on both surface, glandular hairs denser on lower surface. **Inflorescence:** raceme, axillary, open, 5–13flowered, bracteate, puberulent with scattered, subsessile glandular hairs having yellowish heads; peduncle strongly 5-ridged, to 15 mm long; rachis ridged with a principal ridge descending from each pedicel; bractlet subtending pedicel narrowly acuminate, to 1 mm long, puberulent, early-deciduous from persistent base; pedicel arching, 0.5–0.7 mm long. Flower: bisexual, bilateral, pealike (papilionaceous), 1 mm across; nectary inconspicuous on receptacle below ovary; calyx 5-lobed from midpoint, green, puberulent with subsessile glandular hairs; tube funnel-shaped to narrowly bell-shaped, ca. 1.3 mm diameter; lobes heteromorphic, appressed, lower lobe acuminate-linear, to 2.5 mm long, lateral 2 lobes ca. 1 mm long, upper portion deeply 2-lobed, lobes 1-veined, inner (upper) surface glabrous; petals 5, clawed, in range ca. 5 mm long, the claws whitish, limbs pale orange-yellow to light orange-yellow; banner obovate to roundish, ca.  $5 \times 4$  mm, claw blunt, < 1 mm long, limb folded over flower, with a pair of short teeth at base, outer (lower) surface with raised veins and puberulent with subsessile glandular hairs, inner (upper) surface with radiating nectar lines; wings 2, < banner, claws slender, ca. 1.2 mm long, limbs obovate, ca.  $3 \times 1.3$ mm, with side-pointing basal lobe deltate, 0.5 mm long, whitish, glabrous; keel of 2 petals fused on lower side of limbs, = banner, glabrous, claws slender, ca. 1 mm long, whitish, limb broadly elliptic, ca.  $3.5 \times 1.5$  mm; stamens 10, diadelphous (9 fused and 1 free), ca. 5 mm long; filament sheet 2.5–3 mm long, whitish to pale green, free portions alternating long and short, glabrous; anthers dorsifixed, dithecal, some slightly exserted, monomorphic,  $\pm 0.35$  mm long, pale orange-yellow, longitudinally dehiscent; pollen pale yellow; **pistil** 1, ca. 5 mm long, slightly exserted; ovary superior, ellipsoid, to 2.5 mm long, light green, short-sericeous with upward-pointing hairs, 1-chambered with 2–3 ovules attached on upper side; style cylindric, pale green aging white; stigma terminal, capitate, papillate. Fruit: pod (legume), dehiscent with valves slightly twisting (to 180°), mostly 2seeded, low hatchet-shaped and flat but bulging at seeds,  $14-21 \times 2.5-4$  mm, brown to dark brown, with short point at tip, short-pilose and gland-dotted with ruby red heads. **Seed:** plump kidney-shaped,  $3.2-3.5 \times 2-2.2$  mm, hard, greenish to brown mottled usually black or light gray, glossy or not; hilum median. A. C. Gibson